

## N,N'-(1S)-[1,1'-Binaphthalene]-2,2'-diylbis-(2S,2'S)-pyrrolidine-2-carboxamide

Gabriela Guillena, University de Alicante, Alicante, Spain

Carmen Nájera, University de Alicante, Alicante, Spain

*Encyclopedia of Reagents for Organic Synthesis*


Copyright © 2008 John Wiley & Sons, Ltd

[DOI](https://doi.org/10.1002/047084289X.rn01044): 10.1002/047084289X.rn01044

Article Online Posting Date: September 15, 2009

[Previous](#)

### Abstract



[891496-34-7] C<sub>30</sub>H<sub>30</sub>N<sub>4</sub>O<sub>2</sub> (MW 478.24)

InChI = [1/C30H30N4O2/c35-29\(25-11-5-17-31-25\)33-23-15-13-19-7-1-3-9-21\(19\)27\(23\)28-22-10-4-2-8-20\(22\)14-16-24\(28\)34-30\(36\)26-12-6-18-32-26/h1-4,7-10,13-16,25-26,31-32H,5-6,11-12,17-18H2,\(H,33,35\)\(H,34,36\)/f/h33-34H](#)

InChIKey = [FYTHJMKJVUTERE-UBXIPSODCL](#)

(used as a catalyst and as a ligand in enantioselective synthesis of aldol products and cyanosilylation of ketones, respectively)

*Alternate Names:* (S<sub>a</sub>)-2,2' -bis[[(2S)-pyrrolidin-2-ylcarbonyl]amino]-1,1' -binaphthalene, (S)-Binam-L-Pro.

*Physical Data:* mp 230 °C;  $[\alpha]_{26}^D = -108.6^\circ$  (c = 1, MeOH).

*Solubility:* soluble in most polar organic solvents; insoluble in hexane, ether, and

water.

*Form Supplied in:* not commercially available. Prepared from (*S<sub>a</sub>*)-2,2' -diamino-1,1' -binaphthalene and protected l-proline, both commercially available.

*Analysis of Reagent Purity:* elemental analysis, NMR, HRMS, and X-ray structure.


*Preparative Methods:* the reagent can be prepared by coupling (*S<sub>a</sub>*)-2,2' -diamino-1,1' -binaphthalene with the mixed anhydride obtained from l-Boc-proline and isobutyl or ethyl chloroformate or by direct coupling of Boc-L-proline with (*S<sub>a</sub>*)-2,2' -diamino-1,1' -binaphthalene (Binam) mediated by 1-ethyl-3-(3-methylaminopropyl)carbodiimide (EDC) and 1-hydroxybenzotriazole (HOBt) or *N,N'* -dicyclohexylcarbodiimide (DCC) and 4-(dimethylamino)pyridine (DMAP). The coupled intermediate was deprotected by using TFA. Alternatively, the reagent can be prepared by coupling (*S<sub>a</sub>*)-2,2' -diamino-1,1' -binaphthalene with the acyl chloride derived from Fmoc-l-proline, and deprotection with piperidine.

*Purification:* by column chromatography and/or recrystallization from CH<sub>2</sub>Cl<sub>2</sub>/hexane or CHCl<sub>3</sub>/ether.

*Handling, Storage, and Precaution:* the reagent is stable for months when stored in air at room temperature.

#### Enantioselective Cyanosilylation of Ketones


The generation of chiral quaternary stereocenters is a challenging task, with the addition of trimethylsilyl cyanide (TMSCN) to ketones being one of the most successful examples.<sup>1</sup> (*S*)-Binam-l-Pro was used as a ligand for the enantioselective addition of TMSCN to acetophenone (eq 1). The catalytic complex obtained by reaction with titanium tetra isopropoxide gave the O-TMS cyanohydrin with 47% yield and 33% ee. Better results (up to 90% yield and 94% ee) were obtained under similar reaction conditions using the bisamide derived from 1,2-diphenylethane-1,2-diamine and proline.<sup>2</sup>




#### Enantioselective Direct Aldol Reaction

The use of organocatalytic methods<sup>3</sup> for the enantioselective direct aldol reaction<sup>4</sup>


has reached its maturity allowing the synthesis of chiral molecules with high atom efficiency.<sup>5</sup> (S)-Binam-L-Pro was first used as an organocatalyst in the intermolecular direct aldol reaction of aromatic aldehydes and alkyl or cyclic ketones using two different protocols.<sup>6</sup> Whereas in one protocol the mixture of 1,4-dioxane/ketone (4:1) at 4 °C were the best reaction conditions, giving the corresponding aldols with 9-79% yield and 50-88% ee's (eq 2),<sup>6a</sup> DMF/water (1:1) at 0 °C or DMF at 25 °C afforded better results in the second instance (52-99% yield, 78-95% ee).<sup>6b</sup> Under the latter reaction conditions, the use of l-proline as catalyst afforded the racemic product.




When 2-butanone was used as precursor, the reaction took place nearly exclusively at the methyl position, to give the *iso*-isomer preferentially (eq 3). These conditions permitted the recovery by acidic-basic extraction and reuse of catalyst without any detrimental effect on the obtained yields and enantioselectivities during three additional cycles.<sup>6b</sup> Alternatively,  $\text{CHCl}_3$ /ketone (1:1) at -27 °C gave the corresponding aldols with 35-98% yield and 68-95% ee.<sup>7</sup>




Under similar conditions, (S)-Binam-l-Pro was used as catalyst in the reaction between  $\alpha$ -alkoxy ketones ( $R^1 = \text{OH}, \text{OMe}, \text{OBn}, \text{OTBDMS}$ ) and aromatic aldehydes (eq 4) to give mainly the *anti*/*syn*-regioisomer mixture, with small amounts of corresponding *iso*-regioisomer being formed. The diastereoselectivity was dependent on the nature of  $R^1$  group, with the *anti*-regioisomer obtained as the main product. The enantioselectivity of the process ranged from 73 to 99%. For the case of  $\alpha$ -hydroxyacetone ( $R^1 = \text{OH}$ ), the best conditions employed DMSO at 25 °C, affording mainly the *anti*-isomer with 85% ee. These results are comparable in terms of regio- and diastereoselectivities to those obtained with l-proline, with the advantage that (S)-Binam-l-Pro prolinamide could be recovered when DMF was used as solvent.<sup>8</sup>





The reaction rate was highly increased by the addition of catalytic amounts of carboxylic acids, with benzoic acid giving the best results. For instance, using 20 mol % benzoic acid in the reaction between acyclic and cyclic alkyl ketones with 4-nitrobenzaldehyde in DMF:H<sub>2</sub>O, the reaction time was reduced from 3 d to only 1.5 h, maintaining the enantioselectivity. This procedure enabled reactions at -20 °C with enhanced enantioselectivity (86–99%). Using benzoic acid as cocatalyst allowed the reaction to be conducted in water without addition of DMF.<sup>9</sup> The combination of (S)-Binam-l-Pro (10 mol %) and benzoic acid (20 mol %) in either DMF or pure water permitted the use of less reactive ketones such as  $\alpha$ -(methylsulfanyl)acetone,

giving mainly the *iso*-isomer with an excellent 93% ee (eq 5).<sup>10</sup>   $\alpha$ -Alkoxy ketones give similar yields, regio-, diastereo-, and enantioselectivities to those achieved in the absence of acid, but in shorter reaction times (3-24 h).




Other carboxylic acids can be used as cocatalyst. For instance, using acetic acid (10 mol %) as cocatalyst in toluene at -40 °C, the aldol products were obtained in 45-91% yield, 40-96% de, and 67-95% ee but longer reaction times being required (2-3 d).<sup>11</sup> Using a micellar agent stearic acid (20 mol %) as cocatalyst in water at 2 °C permitted reduction of the amount of nucleophilic ketone to 3 equiv, providing 61-99% yield and 58-93% ee in only 12 h.<sup>12</sup>

Using benzoic acid as cocatalyst,   $\alpha$ -chloroacetone mainly produces the *anti*-isomer in moderate yields, but high diastereo- and enantioselectivities. These aldol products were easily converted into chiral (3*R*,4*S*)-*trans*-epoxides by treatment with triethylamine (eq 6).<sup>13</sup>




A further improvement involved the use of solvent-free conditions. Thus, mixing the reagents by simple magnetic stirring allowed reduction of nucleophile to only 2 equiv, catalyst loading of 5 mol %, and in some cases reduced reaction time. The aldol reaction between cycloalkyl, alkyl, and  $\alpha$ -functionalized ketones with aldehydes gave the expected aldol products with similar yields, regio-, diastereo-, and enantioselectivities to those obtained in solution. Furthermore, aldehydes can also be used as nucleophiles providing, after in situ reduction of the aldol products, chiral 1,3-diols with moderate to good enantioselectivities mainly as *anti*-isomers (eq 7).<sup>14</sup>












### Enantioselective Mannich Addition

A single example of the use of (*S*)-Binam-L-Pro (10 mol %) as catalyst for this type of transformation has been reported.<sup>7</sup> Thus, the multicomponent<sup>15</sup> Mannich reaction between 4-nitrobenzaldehyde, 4-methoxyaniline, and acetone afforded the corresponding  $\beta$ -amino ketone with 51% ee at 25 °C (eq 8).



## Bibliography

1. (a) Ramón, D. J.; Yus, M., *Curr. Org. Chem.*, 2004, **8**, 149. [Links](#) (b) *Quaternary Stereocenters: Challenges and Solutions for Organic Synthesis*; Christoffers, J.; Baro, A., Eds.; Wiley-VCH Verlag: Weinheim, 2005. (c) Ramón, D. J.; Yus, M., *Chem. Rev.* 2006, **106**, 2126. [Links](#)
2. Xiong, Y.; Huang, X.; Gou, S.; Huang, J.; Wen, Y.; Feng, X., *Adv. Synth. Catal.* 2006, **348**, 538. [Links](#)
3. (a) Berkessel, A.; Gröger, H., *Asymmetric Organocatalysis: From Biomimetic Concepts to Application in Asymmetric Synthesis*; Wiley-VCH: Weinheim, 2005. (b) *Enantioselective Organocatalysis*; Dalko, P. I. Ed.; Wiley-VCH: Weinheim, 2007. (c) Guillena, G.; Ramón, D. J., *Tetrahedron: Asymmetry* 2006, **17**, 1465. [Links](#) (d) Guillena, G.; Ramón, D. J.; Yus, M., *Tetrahedron: Asymmetry* 2007, **18**, 693. [Links](#)
4. Guillena, G.; Nájera, C.; Ramón, D. J., *Tetrahedron: Asymmetry* 2007, **18**, 2249. [Links](#)
5. (a) Trost, B. M., *Science* 1991, **254**, 1471. [Links](#) (b) Sheldon, R. A., *Pure Appl. Chem.* 2000, **72**, 1233. [Links](#) (c) Trost, B. M., *Acc. Chem. Res.* 2002, **35**, 695. [Links](#) (d) Trost, B. M.; Frederiksen, M. U.; Mathias, U.; Rudd, M. T., *Angew. Chem., Int. Ed.* 2005, **44**, 6630. [Links](#) (e) Guillena, G.; Ramón, D. J.; Yus, M., *Angew. Chem., Int. Ed.* 2007, **47**, 2358. [Links](#)
6. (a) Gryko, D.; Kowalczyk, B.; Zawadzki, L., *Synlett* 2006, 1059. [Links](#) (b) Guillena, G.; Hita, M. C.; Nájera, C., *Tetrahedron: Asymmetry* 2006, **17**, 729. [Links](#)
7. Guizzetti, S.; Benaglia, M.; Pignataro, L.; Puglisi, A., *Tetrahedron: Asymmetry* 2006, **17**, 2754. [Links](#)
8. Guillena, G.; Hita, M. C.; Nájera, C., *Tetrahedron: Asymmetry* 2006, **17**, 1027. [Links](#) (Corrigendum: *Tetrahedron: Asymmetry* 2007, **18**, 1030). [Links](#)
9. Guillena, G.; Hita, M. C.; Nájera, C., *Tetrahedron: Asymmetry* 2006, **17**, 1493. [Links](#)

- [Links](#) (Corrigendum: Tetrahedron: Asymmetry 2007, **18**, 1031).  [Links](#)
10. Guillena, G.; Hita, MC.; Nájera, C.; ARKIVOC 2007, iv, 260.  [Links](#)  
(Corrigendum: ARKIVOC 2007, i, 146).  [Links](#)
11. Ma, G. N.; Zhang, Y. P.; Shi, M., Synthesis 2007, 197.  [Links](#)
12. Guizzetti, S.; Benaglia, M.; Raimondi, L.; Celentano, G., Org. Lett. 2007, **9**, 1247.  [Links](#)
13. Guillena, G.; Hita, M. C.; Nájera, C.; Tetrahedron: Asymmetry 2007, **18**, 1272.  [Links](#)
14. (a) Guillena, G.; Hita, M. C.; Nájera, C.; Vióquez, S. F., Tetrahedron: Asymmetry 2007, **18**, 2300.  [Links](#) (b) Guillena, G.; Hita, M. C.; Nájera, C.; Vióquez, S. F., J. Org. Chem. 2008, **73**, 5933.  [Links](#)
15. Ramón, D. J.; Yus, M., Angew. Chem., Int. Ed. 2005, **44**, 1602.  [Links](#)